



Main Line Health[®]

Well ahead.[®]

Opioid Pathway and Withdrawal Risk Screen

February 21, 2019

OPIOID Withdrawal Risk Screen Launched January 8, 2019

For BMH, LMC, PH, RH

- MLH implemented an Opioid withdrawal risk screening tool to improve identification of patients who are at risk for potential opioid withdrawal.
- Nursing will complete screening questions in EPIC Triage Navigator and ED Narrator (Includes OB and Peds 14yo and older)
- Consist of 2 screening questions:
 - In the past week, have you excessively or inappropriately used prescription opioid pain medication or used heroin.
 - Do you get sick if you cannot use opioid pain medications (including methadone) or heroin?
- Positive screen (yes to either questions) Launches Physician and Nursing BPAs – ED and Inpatient
 - Nursing BPA includes notification to Physician of any positive screen

Nursing Documentation – ED and Inpatient

OVERVIEW

- Allergies
- Pregnancy/Lactation
- Travel/Exposure
- Height/Weight
- Home Meds
- Immunizations
- Vaccinations
- ACP Resources
- Directives
- Filed Documents
- Consents
- Expected Discharge

HISTORY

- Care Everywhere
- Problem List
- Medical/Surgical
- Implants
- LDAs
- AUDIT-C
- Alcohol Use
- Opioid Withdrawal...
- Drug/Tobacco Use

ASSESSMENTS

Opioid Withdrawal Screening - Opioid Withdrawal Risk Screening

Time taken: 1320 12/10/2018

Values By + Create Note


▼ Opioid Withdrawal Risk Screening

In the past week, have you excessively or inappropriately used prescription opioid pain medication or used heroin?

Do you get sick if you cannot use opioid pain medications (including methadone) or heroin?

Provider Notified of positive screen for risk of opioid withdrawal

Notify Provider if YES to either opioid withdrawal risk screening question.



Provider BPA – ED and Inpatient

- Will get BPA if “Yes” to opioid withdrawal screening
- UDS/pregnancy screen will appear if resulted at time of BPA
- Positive screen identifies that the patient is at risk for opioid withdrawal. This does not determine Opioid use disorder, although physician should assess for this.
- Midlevel can act on BPA

BestPractice Advisory - Treatmentplan,Testing

High Priority (1)

BPA ALERT - This patient has been identified to be at risk for opioid withdrawal.

Consider:

- **Urine Drug Screen with Methadone should be ordered on all patients unless already ordered.** (Supplemental Methadone testing can be ordered as *Methadone, Urine* or as a *Supplemental drug screen, urine* which includes Methadone, Methaqualone, and Oxycodone)
- **Consult Social Work and provide "Warm Handoff"**
- **Consult Psychiatry/Telepsychiatry** (All inpatients identified at risk for opioid withdrawal with Urine Drug Screen positive for benzodiazepines and opiates should receive a consult to Psychiatry.)
- **Urine Pregnancy for all appropriate patients.**

Results - last 7 days

Lab	12/18/18 1350	12/18/18 1349
PCP Scrn, Ur	Positive*	--
Benzodiazepine Ur Qual	Positive*	--
Cocaine Screen, Urine	Positive*	--
Amphetamine+Methamphetamine Screen, Ur	Positive*	--
Cannabinoid Screen, Urine	Positive*	--
Opiate Scrn, Ur	Positive*	--
Barbiturate Screen, Ur	Positive*	--
Methaqualone	Positive*	--
Oxycodone Screen, Ur	Positive*	--
Preg Test, Ur	--	Positive*
BHCG, Qualitative Urine	Positive*	--
POCT BhCG, Urine Qual	--	Positive*

Order	Do Not Order	Urine Drug Screen Panel with Methadone
Order	Do Not Order	Consult to Social Work
Order	Do Not Order	Notify physician if patient: SBP<90, HR<60, RR<10, is oversedated.
Order	Do Not Order	Inpatient consult to Psychiatry
Order	Do Not Order	Consult to Telepsychiatry (ED)
Order	Do Not Order	BhCG, Urine, Qual

Acknowledge Reason